

NAME

catanh, catanhf, catanhl - complex arc tangents hyperbolic

SYNOPSIS

```
#include <complex.h>
```

```
double complex catanh(double complex z);
```

```
float complex catanhf(float complex z);
```

```
long double complex catanhl(long double complex z);
```

Link with *-lm*.

DESCRIPTION

The **catanh()** function calculates the complex arc hyperbolic tangent of *z*. If $y = c \operatorname{atanh}(z)$, then $z = c \operatorname{atanh}(y)$. The imaginary part of *y* is chosen in the interval $[-\pi/2, \pi/2]$.

One has:

$$\operatorname{catanh}(z) = 0.5 * (\operatorname{clog}(1 + z) - \operatorname{clog}(1 - z))$$

VERSIONS

These functions first appeared in glibc in version 2.1.

CONFORMING TO

C99.

EXAMPLE

```
/* Link with -lm */
#include <complex.h>
#include <stdlib.h>
#include <unistd.h>
#include <stdio.h>

int
main(int argc, char *argv[])
{
    double complex z, c, f;

    if (argc != 3) {
        fprintf(stderr, Usage: %s <real> <imag>n, argv[0]);
        exit(EXIT_FAILURE);
    }

    z = atof(argv[1]) + atof(argv[2]) * I;

    c = catanh(z);
    printf(catanh() = %6.3f %6.3f*in, creal(c), cimag(c));

    f = 0.5 * (clog(1 + z) - clog(1 - z));
    printf(formula = %6.3f %6.3f*in, creal(f2), cimag(f2));

    exit(EXIT_SUCCESS);
}
```

SEE ALSO

[atanh\(3\)](#), [cabs\(3\)](#), [cimag\(3\)](#), [ctanh\(3\)](#), [complex\(7\)](#)

COLOPHON

This page is part of release 3.74 of the Linux *man-pages* project. A description of the project, information about reporting bugs, and the latest version of this page, can be found at <http://www.kernel.org/doc/man-pages/>.